



# Diagnosis, prognostic factors, and assessment of ALL in adults: 2024 ELN recommendations from a European expert panel

Nicola Gökbüget,<sup>1</sup> Nicolas Boissel,<sup>2</sup> Sabina Chiaretti,<sup>3</sup> Hervé Dombret,<sup>4</sup> Michael Doubek,<sup>5</sup> Adele Fielding,<sup>6</sup> Robin Foà,<sup>3</sup> Sebastian Giebel,<sup>7</sup> Dieter Hoelzer,<sup>1</sup> Mathilde Hunault,<sup>8</sup> David I. Marks,<sup>9</sup> Giovanni Martinelli,<sup>10</sup> Oliver Ottmann,<sup>11</sup> Anita Rijnveld,<sup>12</sup> Philippe Rousselot,<sup>13</sup> Josep Ribera,<sup>14</sup> and Renato Bassan<sup>15</sup>

<sup>1</sup>Department of Medicine II, Hematology/Oncology, Goethe University, University Hospital, Frankfurt, Germany; <sup>2</sup>Hospital Saint-Louis, Assistance Publique–Hôpitaux de Paris, Paris, France; <sup>3</sup>Hematology, Department of Translational and Precision Medicine, Sapienza University, Rome, Italy; <sup>4</sup>Leukemia Department, University Hospital Saint-Louis, Assistance Publique–Hôpitaux de Paris, Saint-Louis Research Institute, Université Paris Cité, Paris, France; <sup>5</sup>Department of Internal Medicine–Hematology and Oncology, University Hospital Brno, Brno, Czech Republic; <sup>6</sup>UCL Cancer Institute, London, United Kingdom; <sup>7</sup>Department of Bone Marrow Transplantation and Onco-Hematology, Maria Skłodowska-Curie National Research Institute of Oncology, Gliwice Branch, Gliwice, Poland; <sup>8</sup>Maladies du Sang University Hospital of Angers, FHU Goal, INSERM, National Centre for Scientific Research, Angers, France; <sup>9</sup>University Hospitals Bristol NHS Foundation Trust, Bristol, United Kingdom; <sup>10</sup>IRCCS Istituto Romagnolo per lo Studio dei Tumori Dino Amadori, Meldola, Italy; <sup>11</sup>Division of Cancer and Genetics, Cardiff University School of Medicine, Cardiff, United Kingdom; <sup>12</sup>Erasmus MC Cancer Institute, Rotterdam, The Netherlands; <sup>13</sup>Clinical Hematology Department, Centre Hospitalier de Versailles, Université Paris-Saclay, Versailles, France; <sup>14</sup>Clinical Hematology Department, Institut Català d'Oncologia-Hospital Germans Trias I Pujol, Josep Carreras Research Institute, Badalona, Spain; and <sup>15</sup>Division of Hematology, Ospedale dell'Angelo, Mestre-Venice, Italy

**Working groups of the European LeukemiaNet have published several important consensus guidelines. Acute lymphoblastic leukemia (ALL) has many different clinical and biological subgroups and the knowledge on disease biology and therapeutic options is increasing exponentially. The European Working Group for Adult ALL has therefore summarized the current state of the art and provided comprehensive consensus recommendations for diagnostic**

**approaches, biologic and clinical characterization, prognostic factors, and risk stratification as well as definitions of endpoints and outcomes. Aspects of treatment, management of subgroups and specific situations, aftercare, and supportive care are covered in a separate publication. The present recommendation intends to provide guidance for the initial management of adult patients with ALL and to define principles as a basis for future collaborative research.**

## Introduction

Scientific progress in acute lymphoblastic leukemia (ALL) has been driven by the activities of national academic multicenter study groups in Europe and US in the past decades. The European national study groups have founded a collaboration within the European LeukemiaNet named European Working Group for Adult ALL. The group has published a consensus recommendation for the management of adult ALL,<sup>1</sup> which was broadly distributed among the participating centers.

ALL is rare compared to other cancers of adulthood. It has many different clinical and biological subgroups and the knowledge on disease biology and therapeutic options is increasing exponentially. There is an unmet need for guidance for clinicians, regulatory agencies, and health care systems. Available guidelines from the National Comprehensive Cancer Center<sup>2</sup> or European Society of Medical Oncology<sup>3</sup> are relevant but need to be complemented by a broad and comprehensive expert consensus considering the different regulatory and socio-economic framework for ALL management in Europe. Therefore, the group decided to develop

European LeukemiaNet Recommendations as published for other entities.<sup>4,5</sup> The focus of this article is on diagnostic approaches, biological and clinical characterization, risk stratification, and definitions of endpoints and outcomes. Aspects of treatment, management of subgroups and specific situations, aftercare, and supportive care are covered in a separate publication.<sup>6</sup>

## Methods

The panel includes 17 members representing national study groups. Members met in person and defined topics, tables, and responsibilities of coauthors (supplemental Table 1, available on the *Blood* website). Coauthors performed literature searches of PubMed database and considered relevant abstracts. The manuscript was reviewed by all coauthors. Formal corrections were performed by the corresponding author. Disagreements were summarized and discussed in the whole group. The whole group agreed on the final version of the manuscript. Due to rapid innovation and availability of new data together with a lack of randomized trials for many essential questions, most of the statements have an evidence level of “expert recommendation” for clinical practice.

## Diagnostic procedures and classification

The initial workup should be comprehensive to allow a precise diagnostic characterization and an accurate stratification, as well as to set the basis for minimal residual disease (MRD) monitoring. Morphology, multicolor flow cytometry (MFC), and molecular genetics are obligatory (Table 1). Additional genomic analyses are used to better define molecular subgroups. Diagnosis is usually based on a bone marrow (BM) aspirate, which should be attempted even in cases with high peripheral blast (PB) counts. A BM biopsy is recommended in patients with a dry aspirate; in these cases, BM aspiration can be repeated after the prephase to obtain further material. Biobanking is recommended. It should include nucleic acid (DNA and RNA), viable cells, and possibly germline material. Further workup includes a lumbar puncture with classification and imaging procedures as needed to define potential extramedullary involvement (supplemental Table 2).

### Morphology and cytochemistry

BM morphology is necessary to evaluate the degree of infiltration and to differentiate ALL from acute myeloid leukemia (AML) and ALL from lymphoblastic lymphoma. There are no

specific cytochemistry reactions to classify ALL. Myeloperoxidase is always negative with the exception of mixed-phenotype acute leukemia (MPAL), which may show low/dim/strong levels of expression particularly in B-myeloid cases in which isolated myeloperoxidase (MPO) expression (isoMPO) may occur.

### Immunophenotyping

MFC (at least 8 colors) is central. It allows to (1) make a differential diagnosis with AML, (2) establish lineage affiliation and differentiation, (3) define an aberrant phenotype for MRD monitoring, and (4) detect target antigens for immunotherapy. The EuroFlow Consortium has standardized MFC.<sup>7</sup> TdT is expressed by all B- and T-cell progenitors, except for mature B-ALL (Burkitt leukemia), while being negative in AML.

B-lineage ALL (B-LIN) accounts for 75% to 80% of cases. The crucial markers are CD19, CD22, and cytoplasmic (cy) CD79a. The EGIL classification<sup>8</sup> defines 4 differentiation stages: pro-B, common, pre-B ALL, and mature B-ALL (Table 1). An aberrant coexpression of myeloid markers can be detected in about 40%<sup>9</sup> of cases. Pro-B ALL is CD10-negative and often

**Table 1. Specific diagnostic workup in adult ALL**

	Recommendation	Outputs
Morphology	Mandatory	Defines infiltration (>25% required for a diagnosis of ALL vs lymphoblastic lymphoma) Allows a differential diagnosis with AML (MPO <sup>-</sup> vs + MPO <sup>+</sup> ) Recognizes L3 (Burkitt-type) subsets
Flow cytometry	Mandatory	Allows a differential diagnosis with AML Permits to define the cell of origin Permits to define the stage of differentiation
MPO*	Mandatory	Allows a differential diagnosis with AML.
B-lineage ALL*: CD19, cCD79a, c/sCD22 (minimal requirement), TdT, CD10, CD20, clgM, slg (kappa or lambda)	Mandatory	Pro-B: CD19/CD79a/cCD22 <sup>+</sup> /CD10 <sup>-</sup> (B-I), NG2 Common: CD10 <sup>+</sup> /clgM <sup>-</sup> (B-II) Pre-B: clgM <sup>+</sup> /slg <sup>-</sup> (B-III) Mature: slg <sup>+</sup> (B-IV), CD20 <sup>+</sup>
T-lineage ALL*: c/sCD3, CD7 (minimal requirement), TdT, CD1a, CD2, CD5, CD4, CD8, TCR $\alpha/\beta$ or $\gamma/\delta$	Mandatory	Pro-T: cCD3/CD7 <sup>+</sup> (T-I) Pre-T: CD2/CD5 <sup>+</sup> (T-II) Cortical-T: CD1a <sup>+</sup> (T-III) Mature-T: sCD3 <sup>+</sup> /CD1a <sup>-</sup> (T-IV)
Molecular genetics	Mandatory	t(9;22)/Ph <sup>+</sup> /BCR::ABL1 t(4;11) <sup>+</sup> /KMT2Ar t(1;19) <sup>+</sup> /TCF3::PBX1 Other high-risk cytogenetics
MRD (molecular or MFC)	Mandatory	Individual MRD assay for further follow-up
Extended genomics (GEP, CNA, WES, WGS, NGS)	In clinical trials and for research purposes	Identification of novel subgroups with prognostic/biologic significance Identification of molecular targets for targeted therapies

c, cytoplasmic; GEP, gene expression profiling; MPO, myeloperoxidase; s, surface; WES, whole exome sequencing; WGS, whole genome sequencing.

\*MPAL: myeloid myeloperoxidase or at least 2 of the following antigens: nonspecific esterase, CD11c, CD14, CD64, and lysozyme; B-lineage: strong CD19 expression plus at least 1 of the following strongly expressed antigens: CD79a, cyCD22, CD10, or weak CD19, with at least 2 of the following strongly expressed antigens: CD79a, cyCD22, CD10; and T-lineage: cCD3 or rarely sCD3.

associated with t(4;11)/KMT2A,<sup>10</sup> whereas pre-B ALL often carries t(1;19)/TCF3::PBX1 aberration. The identification of surface markers as potential targets for immunotherapy is essential for therapeutic approaches.<sup>6</sup>

T-lineage ALL (T-LIN) represents 20% to 25% of adult ALL. Crucial markers are cyCD3 and sCD7. Furthermore, CD1a, CD2, sCD3, CD4, CD5, and CD8, as well as T-cell receptor  $\alpha/\beta$  or  $\gamma/\delta$ , may be variably expressed. T-ALL can be classified into 4 subtypes: pro-T-ALL, pre-T-ALL, cortical (thymic) T-ALL, and mature T-ALL.<sup>8</sup> Pro-T-ALL cases express only cyCD3 and CD7, cortical T-ALL is CD1a positive, whereas mature T-ALL express CD4 and/or CD8, and usually sCD3. The so-called ETP-ALL (early-T precursor),<sup>11</sup> represents an entity within pro-T/pre-T ALL with weak CD5 expression which shows at least one myeloid and/or stem cell marker.<sup>11</sup> CD34 and myeloid markers are expressed in a proportion of T-ALL.<sup>9</sup> The maturation stage correlates with molecular aberrations such as LMO2/HOXA in immature, TLX1/TLX3 in cortical, and TAL1 in mature T-ALL. Besides EGIL there is no uniform international classification of immunophenotypes. To achieve comparability, it is recommended to report results including this classification.

MPAL represents 1% to 5% of acute leukemias<sup>12</sup> and is characterized by blasts coexpressing antigens of more than one lineage on the same cells or that have separate populations of blasts of different lineages. The definition of MPAL has been refined in the World Health Organization (WHO) 2008<sup>13</sup> and 2016<sup>14</sup> classifications, the latter discriminating B-myeloid from the other MPAL. The combination of antigens used for MPAL recognition is summarized in Table 1.

## Cytogenetics and molecular genetics

Karyotyping, fluorescence in situ hybridization, and reverse transcription polymerase chain reaction techniques are used for the genetic characterization.

**B-LIN** t(9;22)/BCR::ABL1 is the most frequent aberration. Its detection in the shortest possible time is essential.<sup>15</sup> Other subtypes include KMT2A rearrangements, monosomy 7, hypodiploidy/low hypodiploidy (and the related near-triploid), t(1;19)/TCF3::PBX1, and t(17;19)/TCF3::HLF. t(8;14)/MYC/IGH are rarely detected; also t(12;21)/ETV6::RUNX1 and high hypodiploidy (51-65 chromosomes) are rare in adults.<sup>16</sup>

**T-LIN** Genetic aberrations often comprise the 14q11 and 7q34 breakpoints, leading to the juxtaposition of the TCR loci to transcription factors, namely TAL1, TAL2, LYL1, LMO1, LMO2, TLX1 (HOX11), TLX3 (HOX11L2), HOXA, MYC, and MYB.<sup>16</sup> In addition, cryptic ABL-1 rearrangements with different partner genes for example, NUP214, EML1, and ETV6 have been identified.

**MPAL** The list of genomic lesions detected in MPAL is rather large, and is likely to increase, given its heterogeneity. As already mentioned, the most frequent rearrangements are BCR::ABL1 detected in about 15% of cases and KMT2A found in roughly 10% of patients; ZNF384 rearrangements (49% of B/MPAL) and BCL11B-r have also been reported.<sup>17,24</sup>

## Extended genomics

Gene expression profiling, copy number alteration analysis (CNA), genome-wide, and next generation sequencing (NGS) techniques have identified new subgroups. These techniques are not part of the standard workup but may be carried out for a refined classification in research laboratories.

The Philadelphia (Ph)-like subgroup is discussed separately.<sup>6</sup> ETP-ALL<sup>11</sup> is genetically heterogeneous, with mutations of multiple pathways including hematopoietic and lymphoid development, Ras, cytokine receptor, kinase signaling, and loss-of-function mutations targeting epigenetic regulators.<sup>18,19</sup>

CNA allows to identify aberrations that affect cell cycle (CDKN2A/B and RB1), as well as lymphoid development (IKZF1, PAX5, VPBEB1, and LEF1). Among those, IKZF1 deletion ( $\Delta$ IKZF1)<sup>20,21</sup> can be found in about 80% of Ph<sup>+</sup> and Ph-like cases.<sup>22</sup> The IKZF1-plus genotype (ie, iKZF1 plus CDKN2A/B and/or PAX5 deletions) is considered in some trials.

Recently, further subtypes have been identified in B-LIN. The DUX4- and ERG-rearrangement leads to the formation of an ERG isoform, which in mice models induces leukemic transformation. MEF2D rearrangements have transforming capability in vitro. Finally, ZNF384 deregulation, which can rearrange with several partners (EP300, CREBBP, TAF15, SYNRG, EWSR1, TCF3, and ARID1B) often displays a weak CD10 and CD13/CD33 expression.<sup>16</sup>

NGS has identified novel mutations and rearrangements; the most frequent involve the RAS pathway (N/K RAS, FLT3, PTPN11, NF1, and BRAF mutations) and can be detected in more than 40% of cases, are prevalent in the hyperdiploid and KMT2A-rearranged cases and tend to increase at relapse. More rare mutations affect the Janus kinase (JAK)/STAT pathway (JAK1/2, JAK3, IL7R, rarely CRLF2, SH2B3, and IL2RB). RAS and JAK/STAT mutations are detected in both B- and T-LIN.<sup>23</sup> In T-ALL, NOTCH1/FBXW7 lesions represent the most frequent event (60%) (Table 2).

## WHO and ICC

The updated classifications are largely overlapping; the International Consensus Classification (ICC) identifies more distinct molecular categories<sup>16,24</sup> (supplemental Table 3). A variety of methods would be required to establish the respective categories. This is not standard in most countries and beside the most relevant diagnostic characterizations summarized in Table 1 the new subcategories to date have limited immediate clinical consequences. Nevertheless, their identification is of interest in a research context or for potential use for selection of relapse therapy. Further WHO and ICC classifications with more detailed guidance on recommended and standardized methods for identification of subgroups are warranted.

## MRD testing

MRD testing aims at detecting and quantifying residual blasts beyond the sensitivity of cytomorphology, which is around 5%.<sup>25,26</sup> It is important to establish the target structures for MRD testing based on primary diagnostic material. The panel strongly advises to perform MRD evaluation in reference laboratories participating in standardization approaches.

**Table 2. Molecular subgroups in adult ALL: incidence, prognosis, and molecular findings**

ALL subset	Prevalence and prognosis*	Related aberration(s)
<b>B-lineage ALL</b>		
<i>BCR::ABL1</i> /t(9;22)(q34;q11.2) (Ph <sup>+</sup> )	20%-50%, increasing with age; improved by tyrosine kinase inhibitor therapy	<i>BCR::ABL1</i> rearrangement
Ph-like	25%-27% Unfavorable/controversial for current regimens	Gene expression profile like <i>BCR::ABL1</i> <sup>+</sup> ALL but without <i>BCR::ABL1</i> rearrangement
<i>TCF3::PBX1</i> /t(1;19)(q23;p13)	10%-15% Favorable with intensive therapy	<i>TCF3::PBX1</i> rearrangement
<i>KMT2A(MLL)::AFF1</i> /t(4;11)(q21;q23.3), <i>KMT2A</i> -rearranged/t(v;11q23.3)	~5% Unfavorable	<i>KMT2A::AFF1</i> or <i>KMT2A</i> -other partner gene rearrangement
<i>IGH::MYC</i> /t(8;14)(q24;q32)	1%-5% Unfavorable in B-precursor ALL	<i>IGH::MYC</i> rearrangement
<i>TCF3::HLF</i> /t(17;19)(q22;p13.3)	<1% Unfavorable	<i>TCF3::HLF</i> rearrangement
<i>iAMP21</i>	~2% Intermediate/unfavorable	—
14q32 translocations	<5%, higher in adolescents Intermediate/unfavorable	<i>IGH</i> fusion with partner genes <i>CRLF2</i> , <i>ID4</i> , <i>CEBP</i> , <i>BCL2</i> , <i>EPOR</i> , <i>LHX4</i> , and <i>IL-3</i>
9p13 deletions/translocations	~25% No impact on outcome	<i>PAX5</i> fusion with partner genes <i>ETV6</i> , <i>ELN</i> , <i>POM121</i> , <i>PML</i> , <i>FOXP1</i> , <i>MLLT3</i> , <i>JAK2</i> , <i>C20orf112</i> , <i>AUTS2</i> , <i>CHFR</i> , <i>SOX5</i> , and <i>POM121C</i>
7p12.2 focal deletions/mutations	50%; 80% in Ph <sup>+</sup> and Ph-like Controversial prognosis	Deletions of <i>IKZF1</i>
<i>DUX4</i> -rearranged and <i>ERG</i> -deregulated	3%-7% Favorable	<i>ERG</i> and <i>IKZF1</i> deletions
<i>MEF2D</i> -rearranged ALL	3%-4% Poor	
<i>ZNF384</i> -rearranged ALL	6%-7% Intermediate	Partner gene <i>EP300</i> , <i>CREBBP</i> , <i>TAF15</i> , <i>SYNRG</i> , <i>EWSR1</i> , <i>TCF3</i> , and <i>ARID1B</i>
<i>CDX2::UBTF</i>	2.7%-4% Poor	High expression of <i>CDX2</i> and gain (1q); <i>UBTF::ATXN7L3</i> ; <i>CDX2</i> -cis-deregulation
<b>T-lineage ALL</b>		
<i>TAL</i> and <i>LMO</i> rearrangements/del(1)(p32), t(1;14)(p32;q11), t(1;7)(p32;q34), t(7;9)(q34;q32), t(11;14)(p15;q1), t(11;14)(p13;q1), t(7;11)(q35;p13)	30%-40% Favorable and partly depending on additional lesions	<i>SIL-TAL1</i> rearrangement, TCR rearrangements with <i>TAL1</i> , <i>TAL2</i> , <i>LMO1</i> , and <i>LMO2</i>
<i>HOXA</i> aberrations/inv(7)(p15q34), t(7;7)(p15;q34), t(10;11)(p13;q14), t(v;11q23), del(9)(q34)	20%-25% Outcome depending on additional lesions	<i>TCR-HOXA</i> rearrangement, <i>MLLT10</i> and <i>MLL</i> rearrangements with various partners, <i>SET-NUP214</i> rearrangement
<i>TLX1-10q24</i> rearrangements/t(7;10)(q34;q24), t(10;14)(q24;q11)	20%-30% No impact on prognosis	<i>TCR-TLX11</i> rearrangement
<i>ETP</i> ALL	10%-15% Unfavorable/controversial	Deregulation of myeloid transcription factors, of members of RAS pathway and of epigenetic regulators
<i>TLX3-5q35</i> rearrangement/t(5;14)(q35;q32)	10% No impact on prognosis	<i>TLX3-BCL11B</i> rearrangement
t(8;14)(q24;q11)	1% Unfavorable	<i>MYC</i> involvement
<i>ABL1</i> rearrangements	~3% Potentially targetable by tyrosine kinase inhibitors	<i>NUP214</i> , <i>EML1</i> ; <i>ETV6</i>
<i>LYL/MEF2C</i> rearrangement and immature cluster/t(7;19)(q34;p13), del(5)(q14)	3%-17% Unfavorable; improved by intensive treatment	<i>TCR</i> with <i>LYL1</i> <i>MEF2C</i> rearrangements

\*Any prognostic statements should be considered carefully; because they depend on protocol, presence of other prognostic features, and are often based on small patient numbers.

**Table 2 (continued)**

ALL subset	Prevalence and prognosis*	Related aberration(s)
NKX2-1/NKX2-2 rearrangements/ inv(14)(q11.2q13), t(7;14)(q34;q13), inv(14)(q13q32.33), t(14;20)(q11;p11)	6% No impact on prognosis	TCR/IGH-NKX2- or NKX2-2 rearrangements

\*Any prognostic statements should be considered carefully; because they depend on protocol, presence of other prognostic features, and are often based on small patient numbers.

The most common methods for MRD monitoring are MFC and real-time quantitative PCR (RQ-PCR). MFC can be applied to most cases (>90%), can reach a sensitivity of 0.1% to 0.01% ( $10^{-3}$  to  $10^{-4}$ ) and results are promptly available. Disadvantages are as follows: (1) the samples must be rapidly analyzed, (2) the number of evaluable cells is small due to BM hypocellularity, (3) results could be misleading due to phenotypic shift, (4) the interpretation to some extent, is operator dependent, and (5) suboptimal sensitivity. The EuroFlow Consortium has standardized the procedures.<sup>27</sup> Further efforts are ongoing to implement the use of next generation flow mainly for better interoperability and sensitivity.

Molecular MRD monitoring of fusion genes (eg, *BCR::ABL1*) has a sensitivity around 0.01%, is possible in about 40% of cases, is not patient-specific, relatively easy to perform and inexpensive. However, its accuracy is hampered by the variability in the number of RNA transcripts per leukemic cell. The EuroMRD Consortium has standardized MRD detection based on the *BCR::ABL1*.<sup>28</sup>

In most patients, a clonal Immunoglobulin/T-cell receptor (IG/TR) rearrangement can be identified for MRD evaluation. The technology is applicable to over 90% of cases<sup>29</sup> and has been extensively standardized within the EuroMRD Consortium. Technical issues are as follows: (1) in the most immature forms, IG/TR gene rearrangements may not be found; (2) the designed probes may not be sufficiently sensitive; (3) clonal evolution may lead to false negative results; and (4) the amount of diagnostic DNA may be too low.

Any RQ-PCR may fail to precisely define the amount of residual disease in cases with a very low burden, that is, “positive-not-quantifiable”; their identification is an unmet need in clinical practice. These cases are being investigated with novel techniques, such as digital droplet PCR and NGS. For NGS neither methodology nor reporting is standardized. In part, similar technical issues apply as for IG/TR PCR and particularly the claimed higher sensitivity depends on the amount of DNA. MRD testing by NGS is so far not incorporated in the clinical practice in many countries.<sup>30</sup> The various techniques are summarized in supplemental Table 4 and therapeutic application of MRD is discussed separately.<sup>6</sup>

## PFs

Different risk subsets according to disease- and host-related factors can be identified.<sup>31</sup> The individual prognosis must be refined by assessing MRD response<sup>24,25</sup> and its integration with other prognostic factors (PFs). Patients without poor PF and/or with a favorable postinduction MRD course may

represent 50%-60% of all cases and are defined as standard-risk (SR, 5-year overall survival (OS) >50%-60% and up to 70%-80% in selected good-risk subsets), whereas those with any poor PF and/or poor MRD response are classified as high risk (HR, 5-year OS 40% to 50%) (Table 3). This distinction is crucial to develop an effective risk-oriented treatment strategy. One approach is to select HR patients for an allogeneic stem cell transplantation (SCT) and/or other novel therapies, whereas SR patients are treated with chemotherapy with good to very good results and low risk of therapy-related mortality. Risk models depend on treatment protocol, differ among clinical studies and are periodically updated (Table 4).

## Clinical risk factors: age, WBC, and immunophenotype

Older age and high white blood cell counts (WBCs) are universally recognized as predicting poorer outcome.<sup>31</sup> WBC is still a relevant and easily assessable PF. Cutoffs for HR B-ALL are often set  $>30 \times 10^9/L$ . A WBC based stratification in T-ALL ( $>100 \times 10^9/L$ ) is not generally adopted. Other clinical PF are poor performance score (Eastern Cooperative Oncology Group [ECOG] >1), central nervous system–positive ALL in some trials, and undue reductions or delay of postinduction therapy.<sup>32,33</sup> Pro-B-ALL is sometimes defined as HR. Some trials report a poorer prognosis in CD20-positive cases,<sup>34</sup> unless associated with good MRD response<sup>35</sup> or treated with rituximab. In T-ALL the prognosis is worse for the pro/pre-T (or ETP as subentity of immature T-ALL) and mature T subsets, whereas the cortical phenotype has a good prognosis.<sup>36</sup> Outcome of HR T-ALL may be improved using pediatric-based chemotherapy in conjunction with SCT.<sup>37</sup> In newly diagnosed ALL, the BM blast count differentiating ALL from lymphoblastic lymphoma is not considered as a prognostic feature but as a feature identifying a clinically relevant subtype which has an impact on practical management as outlined below. In relapsed ALL BM, blast count can be predictive for response rates.<sup>6</sup>

## Cytogenetics

Recent cytogenetic classifications suffer from lack of information in up to 50% of the patients, low incidence of some aberrations, and interactions with other PFs.<sup>38-40</sup> Good risk karyotypes such as t(12;21) are rare in adults. Most adults fall within an intermediate-risk (IMR) group. According to one of the published classifications, IMR includes the normal diploid subset, hyperdiploidy, and some other abnormalities. Monosomy 7, low hypodiploidy/near triploidy, and complex karyotype with 3-5 simultaneous abnormalities are allocated to a HR cytogenetic category as the t(4;11)/KMT2A rearranged ALL by several groups.<sup>40,41</sup> For complex karyotype, a uniform definition is recommended (Table 3). There is no generally accepted cytogenetic classification for adult ALL and it remains open whether



**Table 3. Potential prognostic and predictive factors in adult ALL**

	Risk factors	Annotations
<b>Patient-related</b>		
Age (y)	>30-60 y (continuous variable) >55 y (older adults and elderly)	Independent PF, usually not affecting risk model (age-adapted protocols)
Performance (ECOG)	>1	Retrospective data; relevance in older patients
<b>Disease-related</b>		
WBC ( $\times 10^9/L$ )	>30 (B), >100 (T)	Variably considered
Immunophenotype	Pro-B, CD20 <sup>+</sup> (B), pro/pre-T, ETP, and mature-T (T)	Variably considered
Cytogenetics and fluorescence in situ hybridization	Ph <sup>+</sup> , t(4;11), hypodiploidy, and complex*	Key prognostic elements; beside Ph <sup>+</sup> and KMT2Ar variably considered
Genetics	<i>BCR::ABL1</i> <sup>+</sup> , <i>KMT2Ar</i> Ph-like, mutated <i>CLRF2/TP53/JAK-STAT</i> , adverse CNA profile (B), unmutated <i>NOTCH1/FBWX7</i> , and abnormal <i>RAS/PTEN</i> (T)	Key prognostic elements Variably considered
Miscellaneous	CNS involvement Poor treatment compliance and undue treatment reductions and delay Pharmacogenomics (affecting antimetabolite disposition) Immune marrow microenvironment Drug response profiling	Occasionally considered Retrospective data, of greater concern with pediatric-type protocols Data in children, not usually assessed in adults Investigational, for research purposes Investigational, for research purposes
<b>Treatment-response dynamics</b>		
Corticosteroid sensitivity (prephase)	Poor prednisone response (PB count $\geq 1 \times 10^9/L$ at the end of prephase)	Historical relevance, occasionally considered
Early/incomplete blast cell clearance (BM morphology)	Day 8-15 or end of induction BM blasts $\geq 5\%$	Variably considered
Time to CR (number of courses)	>1 cycle (late CR)	Variably considered
MRD (molecular/flow cytometry)	MRD positivity (from end of induction onwards): $\geq 0.1\%/0.01\%$ after induction $\geq 0.01\%$ /positive after/during consolidation and pre/post-allogeneic SCT	Key and unifying factor predicting outcome

\*Definition of complex karyotype: 5 or more chromosomal abnormalities excluding those patients with an established translocation.<sup>38</sup>

CNS, central nervous system; ECOG, Eastern Cooperative Oncology Group; and ETP, early thymic precursor.

the prognostic impact of rare aberrations as outlined above is still valid in modern protocols.

### Molecular genetics and genomics

Further genetic aberrations can exert significant prognostic effects (Table 2). Cases with *iAMP21*, Ph-like ALL,<sup>22,42,43</sup> and an altered CNA profile can be associated with a higher relapse risk in both Ph-ALL<sup>44,45</sup> and Ph/*BCR::ABL1*-positive (Ph<sup>+</sup>) ALL.<sup>46,47</sup> An 8-gene CNA profile was validated in pediatric B-ALL.<sup>48</sup> The good-risk CNA classifier consisted of no deletion of *IKZF1*, *CDKN2A/B*, *PAR1*, *EBF1*, and *RB1*; by an isolated deletion of *ETV6*, *PAX5*, and *BTG1*; or by a deletion of *ETV6* with single additional deletion of either *BTG1*, *PAX5*, or *CDK2A/B*. Any other CNA combination exerted a negative prognostic effect. Adult studies identified *KMT2A-AFF1*, Ph-like, low hypodiploid/near haploid, *BCL-2/MYC*-rearranged, *PAX5alt*, *ZNF384r*- or *ZNF384*-like, and *MEF2D*-rearranged as

HR or IMR-HR genotypes.<sup>49,50</sup> *CDX2* deregulated ALL with *UBTF::ATXN7L3* has been described as a new unfavorable subtype,<sup>51-53</sup> which may benefit from targeted therapies.<sup>54</sup>

In T-ALL, overexpression of *HOX11L2* and *ERG*, lack of *NOTCH1* and *FBWX7* mutations, and presence of *RAS* or *PTEN* abnormalities yielded unfavorable outcomes.<sup>55,56</sup> Mutations/alterations of *TP53*, *JAK/STAT*, *BCL-2*, and *MYC* are unrestricted to cell lineage and may confer poor prognosis. As for cytogenetics, it is important to note that prognostic annotations are often based on small patient numbers and not always based on outcomes with current protocols.

### Early response dynamics

Patients with poor blast cell clearance and late responders requiring more than one chemotherapy course to complete remission (CR) have an inferior outcome. The most useful prognostic information for relapse risk and OS and individual

**Table 4. Risk stratification models for allogeneic SCT in adult Ph/BCR::ABL1-negative (Ph<sup>-</sup>) ALL (European study groups)**

National Study Group	Patient age (y)	Risk stratification criteria <sup>a</sup>			
		Postinduction MRD	Cytogenetics/genetics <sup>†</sup>	WBC (×10 <sup>9</sup> /L)	Miscellaneous
GMALL (Germany)	<55	≥0.01% after consolidation (wk 16 onward)	KMT2A <sup>+</sup>	>30 (B)	Late CR, pro-B, early/mature-T
GIMEMA (Italy)	≤65	≥0.01% after early consolidation (wk 10-16), any positivity (wk 22)	Adverse, KMT2A <sup>+</sup>	>100	Early/mature-T
HOVON (The Netherlands)	<40	≥0.01% after consolidation (wk 14-16)	Adverse KMT2A, hypodiploidy, complex karyotype	>30 (B), >100 (T)	Late CR
PALG (Poland)	<55	≥0.1% after induction ≥0.01% during/after consolidation	KMT2A <sup>+</sup>	>30 (B), >100 (T)	CNS <sup>+</sup>
UK NCRI ALL Group (United Kingdom)	<40	≥0.1% after induction and consolidation (mathematical risk model integrating MRD, cytogenetics and WBC)	Adverse	High count	—
FALL (Finland)	<45	≥0.1% after consolidation block B	Abn11q23, hypodiploidy	>100	Late CR, d15 BM blasts >25%
RALL (Russia)	<55	Positive during/after consolidation	t(4;11), t(1;19), KMT2A <sup>+</sup>	—	Age >30
SVALL (Sweden)	<65	≥0.1% after consolidation	Hypodiploidy, KMT2A <sup>+</sup>	—	EOI BM blasts >5%
PETHEMA (Spain)	<55 (60 fit)	≥0.1% after induction ≥0.01% during/after consolidation	—	—	—
GRAALL (France/Belgium/Switzerland)	<60	≥0.1% after induction at wk 6 or ≥0.01% after consolidation at wk 12	—	—	—
CELL (Czech Republic)	<65	≥0.1% after induction ≥0.01% after consolidation	KMT2A <sup>+</sup>	>30 (B)	Early/mature-T

CNS, central nervous system; EOI, end of induction.

<sup>a</sup>Independent risk factors adopted in clinical trials for the definition of HR ALL and the consequent allocation to SCT; adapted from Giebel et al.<sup>63</sup><sup>†</sup>Adverse cytogenetics/genetics (details to be found in single study protocols).

risk stratification comes from MRD testing.<sup>2,25,57-59</sup> As a general rule, with RQ-PCR (sensitivity of 0.01%), good prognosis patients reach MRD levels <0.1% to <0.01% (better if negative) at end of induction (weeks 4-6) and <0.01% (better if negative) following 1 to 3 early consolidation courses around weeks 10-16. Therapeutic consequences are discussed separately.<sup>6</sup>

### Integrated risk models

In children, an independent genotype-specific effect on MRD-related outcomes improved the risk stratification.<sup>60</sup> In adults, one study demonstrated the usefulness of a mixed MRD and genetic risk classification, showing the independent prognostic effect of a 4-gene classifier (favorable: lack of KMT2A rearrangements and ΔIKZF1 in B-ALL; mutated NOTCH1/FWBX7 without RAS/PTEN abnormalities in T-ALL<sup>55</sup>). In addition, a comprehensive risk model combined WBC, cytogenetics, and MRD results.<sup>61</sup> Both models have not been adopted uniformly.

### Recommendations for analysis of PFs and risk stratification

MRD testing is recommended in all cases to optimize risk stratification.<sup>59,62</sup> In a European survey, MRD was the only PF shared by 11 national study groups to define HR ALL and to make a decision about SCT indication<sup>63</sup> (Table 4). No other PF achieved the same level of consensus, although adverse genetics rank high in HR definitions (8/11 groups). Altogether, it is strongly recommended to perform MRD analysis in all patients with ALL.<sup>62</sup> New evidence may prove the independent prognostic power of several genetic abnormalities that in different combinations concur to define novel HR subsets. Appropriate diagnostic identification and clinical data in relevant patient numbers with modern protocols are a prerequisite. New combined risk models are therefore recommended for research purposes to improve the diagnostic and prognostic platforms.

## Response criteria and definition of outcomes

Formal criteria for BM response assessment in ALL are based on AML<sup>4,64</sup> and those for extramedullary response on non-Hodgkin lymphoma (NHL).<sup>65</sup> Recently new standards for remission, treatment failure, and relapse have been defined for pediatric ALL, which should be adopted for adult ALL.<sup>66</sup>

### Cytologic response in BM

Although criteria for CR are uniformly used, CR with incomplete recovery (CRI) can be an endpoint in clinical trials. Failure or partial remission (PR) (Table 5) can be differentiated or combined. For PR cases a correlation with MRD testing is recommended. If technical criteria for MRD testing are met, PR or CRI with negative MRD status can be considered as CR. On the other hand, patients with MRD of 1% or higher can be considered as failure.<sup>66</sup> For future clinical trials, a standardized terminology is requested (Table 5). Publications should also define time point (TP) of response assessment and avoid reporting of cumulative response rates.

### MRD response

Response criteria are redefined by integrating MRD assessment. Usually, MRD assessment is only considered in patients with hematologic CR or CRI. In any case the reference population should be clearly stated and evaluable and nonevaluable MRD results should be named. Complete MRD response is defined as no detection of MRD, with a minimum sensitivity of 0.01% at the respective TP.<sup>29</sup> MRD persistence is defined as quantifiable MRD, usually 0.01% or greater. Some groups consider any MRD positivity as prognostically relevant; others consider thresholds like 0.1% or 0.01% for different TP or different treatment consequences. Beyond these categories additional MRD results may be reported, including nonquantifiable MRD at different levels of sensitivity or negative MRD with insufficient sensitivity.<sup>67,68</sup> For TP with nonquantifiable MRD, additional digital droplet PCR or NGS testing may be used to separate true negative from false positive results.<sup>68</sup> Of note, any negative MRD result without the corresponding sensitivity level at a distinct TP does not fulfill technical requirements. The latter are different for MFC, PCR of IG/TR,<sup>29</sup> PCR of fusion genes,<sup>28</sup> and often not reported for NGS based methods. MRD results should include information on the material tested and specifics: PB vs BM, level, and status of MRD that is, positive vs negative or nonevaluable. Two variants of MRD based response assessment are given in Table 5. The panel is in favor of variant 1 because the exact MRD level (not considered in variant 2) is relevant for any treatment decisions and for comparability of reported results.

For Ph<sup>+</sup> ALL MRD, response criteria are often derived from trials for chronic myeloid leukemia. Whether these categories are meaningful for ALL remains open to discussion. Approaches with an ALL-specific definition are considered. Furthermore, discrepancies between BCR::ABL1 based MRD results and other methods may occur if 2 methods are applied in parallel. BCR::ABL1-based MRD may remain positive whereas IG/TR MR is negative. This is probably due to multilineage involvement with BCR::ABL1 also in myeloid precursors. Clinical consequences are not clear so far, but the ICC has now integrated 2

subgroups with BCR::ABL1 (lymphoid and multilineage).<sup>16</sup> For MRD-based treatment decisions it is recommended to use in addition to BCR::ABL1-based MRD assessment one additional method that is MFC or IG/TR.

### Extramedullary response

Between 20% to 70% of patients with ALL show extramedullary involvement at diagnosis depending on subtype. Type and extent of involvement should be documented. These localizations should have regressed in size at the TP of CR confirmation. Standard criteria for NHL are recommended for classification.<sup>65,69</sup> In case of persistent extramedullary involvement after induction and early consolidation, Positron Emission Tomography (PET) analysis can be considered. This approach raises challenges in dose dense treatment protocols for ALL due to the recommended treatment-free interval. For the purposes of clinical trials, patients with CRI or PR, PET negative may be considered as CR. Therapeutic consequences from positive PET remain to be defined.

### TP for response assessment

The TPs depend on protocol and treatment consequences. Usually, a CR should be achieved after the initial 1 to 3 phases of standard therapy depending on the protocol. Cases not achieving a CR after this period are considered as primary failures. This definition is important for the purpose of clinical trials. During the follow-up, BM response may be assessed every 2 to 3 months until end of maintenance. Further controls may rely on PB. More frequent assessments are recommended in patients with low level MRD of any level or PET positivity at any TP. Any treatment change based on MRD should be followed by MRD assessment of response. Further essential outcome parameters and major approaches for clinical trials are described in the supplement (supplemental Table 5).

## Summary and outlook

The management of ALL is often organized by national study groups and international consortia.<sup>70</sup> Their reference laboratories and/or associated biobanks together with clinical data bases are essential for research on disease biology and prognostication and are sources of reliable real-world data. Reference laboratories are crucial to establish and maintain high standards for diagnostic procedures and biologic characterization as basis for risk stratification and optimal management. Cytomorphology, immunophenotyping, and the identification of prognostically relevant molecular markers together with measurement of MRD are the basis for diagnosis and risk adapted therapy.

Further genetic characterization of samples from biobanks has identified many new subtypes of ALL. The prognostic and therapeutic impact, however, often remains open for current treatment regimens and diagnostic identification is not part of standard of care in many countries. It will be essential to define reliable and cost-effective diagnostic procedures to improve classification of ALL. On this basis, future risk stratification may integrate molecular markers, potential treatment targets, risks for relapse, and toxicities as well as the dynamics of MRD. Furthermore, improved understanding of disease biology can contribute to the development of new and targeted precision medicine approaches.



**Table 5. Response criteria for ALL**

Category	Definition
<b>Hematologic response criteria</b>	
CR*	BM blasts <5% Absence of extramedullary disease Absolute neutrophil count $>1 \times 10^9/L$ Platelet count $>100 \times 10^9/L$ (independence of red cell transfusions) If available: MRD $<1\%$ <sup>66</sup>
CRi†	All CR criteria except for residual thrombocytopenia $<100 \times 10^9/L$ or neutropenia $<1 \times 10^9/L$ If available: MRD $<1\%$ <sup>66</sup>
Morphologic leukemia-free state‡	BM blasts <5% Absence of extramedullary disease If available: MRD $<1\%$ <sup>66</sup>
PR§	Relevant in the setting of phase 1 and 2 clinical trials only; all hematologic criteria of CR; decrease of BM-blast percentage from 5% to 25%; and decrease of pretreatment BM-blast percentage by at least 50% If available: CR if MRD $<1\%$ <sup>66</sup>
Failure	None of the above If available: MRD $\geq 1\%$ <sup>66</sup>
<b>MRD response criteria (variant 1)  </b>	
Complete MRD response	No detectable MRD¶
MRD failure	MRD above 0.01% (ie, $10^{-4}$ )
MRD other	
Negative	MRD negative with insufficient sensitivity
Positive/intermediate	MRD positive below 0.01%, quantifiable MRD positive below 0.01%, nonquantifiable MRD positive, nonquantifiable
<b>MRD response criteria (variant 2)  </b>	
MRD complete response	No detectable MRD¶
MRD persistence	Any quantifiable MRD
<b>Criteria for extramedullary response assessment</b>	Published criteria for NHL <sup>65</sup> PET in case of CRu/PR according to published criteria for NHL <sup>69</sup>

This table is modified from Döhner et al.<sup>71</sup>

\*All criteria need to be fulfilled; marrow evaluation should be based on a count of 200 nucleated cells in an aspirate with spicules; if ambiguous, consider repeat exam after 5 to 7 days; a marrow biopsy should be performed in cases of dry tap, or if no spicules are obtained; no minimum duration of response required.

†CRi is of value in protocols using intensified induction or double induction strategies, in which hematologic recovery is not awaited, but intensive therapy will be continued. In such protocols, CR may even not be achieved during the entire treatment plan. In these instances, the overall remission rate should include CR and CRi patients.

‡This category may be useful in the clinical development of novel agents within phase 1 clinical trials, in which a transient morphologic leukemia-free state may be achieved at the time of early response assessment.

§Marrow should not merely be aplastic; at least 200 cells should be enumerated, or cellularity should be at least 10%.<sup>71</sup> Any PR should be confirmed or falsified by parallel MRD assessment.

||Confirmation of any MRD response requires the application of standardized methods with minimum technical requirements<sup>29</sup> in reference laboratories.

¶Confirmation of negative MRD requires that technical requirements for establishment of sensitivity (usual: 0.01%) of each individual TP are fulfilled.

For comparability of data on an international level it will be essential to use uniform and standardized criteria for reporting of results. This applies particularly for the use of MRD as endpoint of clinical trials and for any MRD-based treatment decisions.

Although there is no standard risk model for adult ALL, all experts agree that depending on treatment protocols and clinical subgroup risk-adapted approaches for management are standard.<sup>6</sup> Future risk models may also integrate

pharmacogenomics to predict and potentially avoid toxicities and in vitro sensitivity testing for identification of potentially active drugs for later-line approaches in relapsed/refractory ALL.

## Authorship

Contribution: All authors reviewed literature, contributed to writing parts of the manuscript, reviewed the manuscript, and approved the final manuscript for submission.

Conflict-of-interest disclosure: N.G. has received institutional research funding from Amgen, Clinigen, Incyte, Jazz, Novartis, Pfizer, and Servier and received speaker honoraria or fees for advisory board participation from Amgen, AstraZeneca, Autolus, Celgene, Clinigen, Gilead, Incyte Jazz, Novartis, Pfizer, and Servier. N.B. received honoraria from Amgen, Incyte, Jazz Pharma, Novartis, Pfizer, and Servier and research grants from Amgen, Novartis, Incyte, Jazz Pharma, and Sanofi. S.C. served on the advisory boards of Incyte, Pfizer, and AbbVie and as a consultant for Gilead and Amgen. H.D. received research funding from Amgen, Astellas, Celgene, Incyte, Jazz, Pfizer, and Servier and honoraria from Daiichi Sankyo, Incyte, Jazz, and Servier. M.D. received research support from AstraZeneca. R.F. received speaker honoraria from Amgen, Novartis, and AstraZeneca and served on the advisory board of Merck Sharp & Dohme. S.G. received speaker honoraria from or joined the advisory boards of Amgen, Novartis, Pfizer, and Gilead; received speaker honoraria from Angelini and travel grants from Gilead; and served on advisory boards of Zentiva. M.H. received research support from Novartis, travel support from AbbVie and BeiGene and speaker honoraria from Servier; joined advisory boards of Servier; and was part of advisory boards of Amgen and Clinigen. D.I.M. provided consultancy for Pfizer, Novartis, and Kite. O.O. received speaker honoraria from, joined advisory boards of, and received research grants from Incyte; received research support from Amgen and Celgene; and received honoraria for advisory boards from Autolus. A.R. received research support from Servier. P.R. received research grants from Pfizer and Incyte. J.R. received speaker honoraria from and joined

advisory boards of Amgen, Pfizer, Shire, Ariad, and Incyte; joined advisory boards of Takeda; and received research support from Amgen. R.B. joined advisory boards of Novartis and Kite/Gilead and received honoraria or travel support from Amgen, Incyte, Servier, Jazz, and Pfizer. The remaining authors declare no competing financial interests.

ORCID profiles: N.G., [0000-0003-2291-8245](https://orcid.org/0000-0003-2291-8245); N.B., [0000-0003-2091-7927](https://orcid.org/0000-0003-2091-7927); M.D., [0000-0002-1269-6282](https://orcid.org/0000-0002-1269-6282); A.F., [0000-0002-4746-7789](https://orcid.org/0000-0002-4746-7789); S.G., [0000-0002-4827-4401](https://orcid.org/0000-0002-4827-4401); M.H., [0000-0001-7777-5216](https://orcid.org/0000-0001-7777-5216); O.O., [0000-0001-9559-1330](https://orcid.org/0000-0001-9559-1330); P.R., [0000-0003-3238-4494](https://orcid.org/0000-0003-3238-4494); J.R., [0000-0003-1042-6024](https://orcid.org/0000-0003-1042-6024); R.B., [0000-0001-8214-2894](https://orcid.org/0000-0001-8214-2894).

Correspondence: Nicola Gökbüget, Department of Medicine II, University Hospital, Theodor Stern Kai 7, 60590 Frankfurt, Germany; email: [goekbuget@em.uni-frankfurt.de](mailto:goekbuget@em.uni-frankfurt.de).

## Footnotes

Submitted 11 July 2023; accepted 21 January 2024; prepublished online on *Blood* First Edition 31 January 2024. <https://doi.org/10.1182/blood.2023020794>.

The online version of this article contains a data supplement.

## REFERENCES

- Gökbüget N, Bassan R, Dombret H, et al, eds. *Recommendations of the European Working Group for Adult ALL*. 1st ed. UNI-MED Verlag AG, D-28323 Bremen. International Medical Publishers; 2011.
- Brown PA, Shah B, Advani A, et al. Acute lymphoblastic leukemia, Version 2.2021, NCCN clinical practice guidelines in oncology. *J Natl Compr Canc Netw*. 2021; 19(9):1079-1109.
- Hoelzer D, Bassan R, Dombret H, et al. Acute lymphoblastic leukaemia in adult patients: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2016;27(suppl 5):v69-v82.
- Dohner H, Estey E, Grimwade D, et al. Diagnosis and management of AML in adults: 2017 ELN recommendations from an international expert panel. *Blood*. 2017; 129(4):424-447.
- Baccarani M, Deininger MW, Rosti G, et al. European LeukemiaNet recommendations for the management of chronic myeloid leukemia: 2013. *Blood*. 2013;122(6): 872-884.
- Gökbüget N, Boissel N, Chiaretti S, et al. Management of ALL in adults: 2024 ELN recommendations from a European expert panel. *Blood*. 2024;143(19):1903-1930.
- Kalina T, Flores-Montero J, van der Velden VH, et al. EuroFlow standardization of flow cytometer instrument settings and immunophenotyping protocols. *Leukemia*. 2012;26(9):1986-2010.
- Bene MC, Nebe T, Bettelheim P, et al. Immunophenotyping of acute leukemia and lymphoproliferative disorders: a consensus proposal of the European LeukemiaNet Work Package 10. *Leukemia*. 2011;25(4): 567-574.
- Vitale A, Guarini A, Ariola C, et al. Absence of prognostic impact of CD13 and/or CD33 antigen expression in adult acute lymphoblastic leukemia. Results of the GIMEMA ALL 0496 trial. *Haematologica*. 2007;92(3):342-348.
- Bueno C, Montes R, Catalina P, Rodriguez R, Menendez P. Insights into the cellular origin and etiology of the infant pro-B acute lymphoblastic leukemia with MLL-AF4 rearrangement. *Leukemia*. 2011;25(3): 400-410.
- Coustan-Smith E, Mullighan CG, Onciu M, et al. Early T-cell precursor leukaemia: a subtype of very high-risk acute lymphoblastic leukaemia. *Lancet Oncol*. 2009;10(2): 147-156.
- Weinberg OK, Dennis J, Zia H, et al. Adult mixed phenotype acute leukemia (MPAL): B/myeloid MPAL<sup>isoMPO</sup> is distinct from other MPAL subtypes. *Int J Lab Hematol*. 2023; 45(2):170-178.
- Vardiman JW, Thiele J, Arber DA, et al. The 2008 revision of the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia: rationale and important changes. *Blood*. 2009;114(5):937-951.
- Arber DA, Orazi A, Hasserjian R, et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood*. 2016; 127(20):2391-2405.
- Foa R, Chiaretti S. Philadelphia chromosome-positive acute lymphoblastic leukemia. *N Engl J Med*. 2022;386(25):2399-2411.
- Arber DA, Orazi A, Hasserjian RP, et al. International Consensus Classification of myeloid neoplasms and acute leukemias: integrating morphologic, clinical, and genomic data. *Blood*. 2022;140(11): 1200-1228.
- Alexander TB, Gu Z, Iacobucci I, et al. The genetic basis and cell of origin of mixed phenotype acute leukaemia. *Nature*. 2018; 562(7727):373-379.
- Zhang J, Ding L, Holmfeldt L, et al. The genetic basis of early T-cell precursor acute lymphoblastic leukaemia. *Nature*. 2012; 481(7380):157-163.
- Neumann M, Heesch S, Schlee C, et al. Whole-exome sequencing in adult ETP-ALL reveals a high rate of DNMT3A mutations. *Blood*. 2013;121(23):4749-4752.
- Mullighan CG, Su X, Zhang J, et al. Deletion of IKZF1 and prognosis in acute lymphoblastic leukemia. *N Engl J Med*. 2009; 360(5):470-480.
- Martinelli G, Iacobucci I, Storlazzi CT, et al. IKZF1 (Ikaro) deletions in BCR-ABL1-positive acute lymphoblastic leukemia are associated with short disease-free survival and high rate of cumulative incidence of relapse: a GIMEMA AL WP report. *J Clin Oncol*. 2009; 27(31):5202-5207.
- Roberts KG, Gu Z, Payne-Turner D, et al. High frequency and poor Outcome of Philadelphia chromosome-like acute lymphoblastic leukemia in adults. *J Clin Oncol*. 2017;35(4): 394-401.
- Mullighan CG. Molecular genetics of B-precursor acute lymphoblastic leukemia. *J Clin Invest*. 2012;122(10):3407-3415.
- Alaggio R, Amador C, Anagnostopoulos I, et al. The 5th edition of the World Health Organization classification of haematolymphoid tumours: lymphoid neoplasms. *Leukemia*. 2022;36(7): 1720-1748.

25. van Dongen JJ, van der Velden VH, Brüggemann M, Orfao A. Minimal residual disease diagnostics in acute lymphoblastic leukemia: need for sensitive, fast, and standardized technologies. *Blood*. 2015; 125(26):3996-4009.
26. Brüggemann M, Kotrova M. Minimal residual disease in adult ALL: technical aspects and implications for correct clinical interpretation. *Blood Adv*. 2017;1(25):2456-2466.
27. Kalina T, Brdickova N, Glier H, et al. Frequent issues and lessons learned from EuroFlow QA. *J Immunol Methods*. 2019; 475:112520.
28. Pfeifer H, Cazzaniga G, van der Velden VHJ, et al. Standardisation and consensus guidelines for minimal residual disease assessment in Philadelphia-positive acute lymphoblastic leukemia (Ph + ALL) by real-time quantitative reverse transcriptase PCR of e1a2 BCR-ABL1. *Leukemia*. 2019; 33(8):1910-1922.
29. Brüggemann M, Schrauder A, Raff T, et al. Standardized MRD quantification in European ALL trials: proceedings of the Second International Symposium on MRD assessment in Kiel, Germany, 18-20 September 2008. *Leukemia*. 2010;24(3):521-535.
30. Brüggemann M, Kotrova M, Knecht H, et al. Standardized next-generation sequencing of immunoglobulin and T-cell receptor gene recombinations for MRD marker identification in acute lymphoblastic leukaemia; a EuroClonality-NGS validation study. *Leukemia*. 2019;33(9):2241-2253.
31. Bassan R, Hoelzer D. Modern therapy of acute lymphoblastic leukemia. *J Clin Oncol*. 2011;29(5):532-543.
32. Advani AS, Jin T, Ramsingh G, et al. Time to post-remission therapy is an independent prognostic factor in adults with acute lymphoblastic leukemia. *Leuk Lymphoma*. 2008;49(8):1560-1566.
33. Kumar AJ, Gimotty PA, Gelfand JM, et al. Delays in postremission chemotherapy for Philadelphia chromosome negative acute lymphoblastic leukemia are associated with inferior outcomes in patients who undergo allogeneic transplant: an analysis from ECOG 2993/MRC UK ALLXII. *Am J Hematol*. 2016; 91(11):1107-1112.
34. Maury S, Huguët F, Leguay T, et al. Adverse prognostic significance of CD20 expression in adults with Philadelphia chromosome-negative B-cell precursor acute lymphoblastic leukemia. *Haematologica*. 2010;95(2):324-328.
35. Mannelli F, Gianfaldoni G, Intermesoli T, et al. CD20 expression has no prognostic role in Philadelphia-negative B-precursor acute lymphoblastic leukemia: new insights from the molecular study of minimal residual disease. *Haematologica*. 2012;97(4):568-571.
36. Goekbuget N, Fiedler W, Alakel N, et al. Results of the risk-adapted, MRD-stratified GMALL trial 08/2013 in 281 T-ALL / T-Lbl patients: excellent outcome of standard risk thymic T-ALL [abstract]. *Blood*. 2022;140(suppl 1):115-117.
37. Bond J, Graux C, Lhermitte L, et al. Early response-based therapy stratification improves survival in adult early thymic precursor acute lymphoblastic leukemia: a group for research on adult acute lymphoblastic leukemia study. *J Clin Oncol*. 2017;35(23):2683-2691.
38. Moorman AV, Harrison CJ, Buck GA, et al. Karyotype is an independent prognostic factor in adult acute lymphoblastic leukemia (ALL): analysis of cytogenetic data from patients treated on the Medical Research Council (MRC) UKALLXII/Eastern Cooperative Oncology Group (ECOG) 2993 trial. *Blood*. 2007; 109(8):3189-3197.
39. Pullarkat V, Slovak ML, Kopecky KJ, Forman SJ, Appelbaum FR. Impact of cytogenetics on the outcome of adult acute lymphoblastic leukemia: results of Southwest Oncology Group 9400 study. *Blood*. 2008; 111(5):2563-2572.
40. Lafage-Pochitaloff M, Baranger L, Hunault M, et al. Impact of cytogenetic abnormalities in adults with Ph-negative B-cell precursor acute lymphoblastic leukemia. *Blood*. 2017; 130(16):1832-1844.
41. Marks DI, Moorman AV, Chilton L, et al. The clinical characteristics, therapy and outcome of 85 adults with acute lymphoblastic leukemia and t(4;11)(q21;q23)/MLL-AFF1 prospectively treated in the UKALLXII/ECOG2993 trial. *Haematologica*. 2013;98(6): 945-952.
42. Jain N, Roberts KG, Jabbour E, et al. Ph-like acute lymphoblastic leukemia: a high-risk subtype in adults. *Blood*. 2017;129(5): 572-581.
43. Herold T, Schneider S, Metzeler KH, et al. Adults with Philadelphia chromosome-like acute lymphoblastic leukemia frequently have IGH-CRLF2 and JAK2 mutations, persistence of minimal residual disease and poor prognosis. *Haematologica*. 2017;102(1): 130-138.
44. Ribera J, Morgades M, Zamora L, et al. Prognostic significance of copy number alterations in adolescent and adult patients with precursor B acute lymphoblastic leukemia enrolled in PETHEMA protocols. *Cancer*. 2015;121(21):3809-3817.
45. Messina M, Chiaretti S, Fedullo AL, et al. Clinical significance of recurrent copy number aberrations in B-lineage acute lymphoblastic leukaemia without recurrent fusion genes across age cohorts. *Br J Haematol*. 2017; 178(4):583-587.
46. Pfeifer H, Raum K, Markovic S, et al. Genomic CDKN2A/2B deletions in adult Ph(+) ALL are adverse despite allogeneic stem cell transplantation. *Blood*. 2018;131(13): 1464-1475.
47. Fedullo AL, Messina M, Elia L, et al. Prognostic implications of additional genomic lesions in adult Philadelphia chromosome-positive acute lymphoblastic leukemia. *Haematologica*. 2019;104(2): 312-318.
48. Hamadeh L, Enshaei A, Schwab C, et al. Validation of the United Kingdom copy-number alteration classifier in 3239 children with B-cell precursor ALL. *Blood Adv*. 2019;3(2):148-157.
49. Mitchell RJ, Kirkwood AA, Barretta E, et al. IKZF1 alterations are not associated with outcome in 498 adults with B-precursor ALL enrolled in the UKALL14 trial. *Blood Adv*. 2021;5(17):3322-3332.
50. Paietta E, Roberts KG, Wang V, et al. Molecular classification improves risk assessment in adult BCR-ABL1-negative B-ALL. *Blood*. 2021;138(11):948-958.
51. Yasuda T, Sanada M, Kawazu M, et al. Two novel high-risk adult B-cell acute lymphoblastic leukemia subtypes with high expression of CDX2 and IDH1/2 mutations. *Blood*. 2022;139(12):1850-1862.
52. Kimura S, Montefiori L, Iacobucci I, et al. Enhancer retargeting of CDX2 and UBTf::ATXN7L3 define a subtype of high-risk B-progenitor acute lymphoblastic leukemia. *Blood*. 2022;139(24):3519-3531.
53. Passet M, Kim R, Gachet S, et al. Concurrent CDX2 cis-deregulation and UBTf::ATXN7L3 fusion define a novel high-risk subtype of B-cell ALL. *Blood*. 2022;139(24): 3505-3518.
54. Bastian L, Hartmann AM, Beder T, et al. UBTf::ATXN7L3 gene fusion defines novel B cell precursor ALL subtype with CDX2 expression and need for intensified treatment. *Leukemia*. 2022;36(6): 1676-1680.
55. Beldjord K, Chevret S, Asnafi V, et al. Oncogenetics and minimal residual disease are independent outcome predictors in adult patients with acute lymphoblastic leukemia. *Blood*. 2014;123(24):3739-3749.
56. Trinquand A, Tanguy-Schmidt A, Ben Abdelali R, et al. Toward a NOTCH1/FBXW7/RAS/PTEN-based oncogenetic risk classification of adult T-cell acute lymphoblastic leukemia: a Group for Research in Adult Acute Lymphoblastic Leukemia study. *J Clin Oncol*. 2013;31(34): 4333-4342.
57. Bassan R, Intermesoli T, Scattolin A, et al. Minimal residual disease assessment and risk-based therapy in acute lymphoblastic leukemia. *Clin Lymphoma Myeloma Leuk*. 2017;17S:S2-S9.
58. Gokbuget N, Kneba M, Raff T, et al. Adult patients with acute lymphoblastic leukemia and molecular failure display a poor prognosis and are candidates for stem cell transplantation and targeted therapies. *Blood*. 2012;120(9):1868-1876.
59. Short NJ, Jabbour E, Albitar M, et al. Recommendations for the assessment and management of measurable residual disease in adults with acute lymphoblastic leukemia: a consensus of North American experts. *Am J Hematol*. 2019;94(2):257-265.
60. O'Connor D, Enshaei A, Bartram J, et al. Genotype-specific minimal residual disease

interpretation improves stratification in pediatric acute lymphoblastic leukemia. *J Clin Oncol*. 2018;36(1):34-43.

61. Moorman A, Kirkwood AA, Enshaeei A, et al. Clinical efficacy of a novel validated prognostic index for trial design in adult acute lymphoblastic leukemia [abstract]. *Hemasphere*. 2019;3(suppl 1):748-749.
62. Gokbuget N. MRD in adult Ph/BCR-ABL-negative ALL: how best to eradicate? *Hematology Am Soc Hematol Educ Program*. 2021;2021(1):718-725.
63. Giebel S, Marks DI, Boissel N, et al. Hematopoietic stem cell transplantation for adults with Philadelphia chromosome-negative acute lymphoblastic leukemia in first remission: a position statement of the European Working Group for Adult Acute Lymphoblastic Leukemia (EWALL) and the Acute Leukemia Working Party of the European Society for Blood and Marrow Transplantation (EBMT). *Bone Marrow Transplant*. 2019;54(6):798-809.
64. Cheson BD, Bennett JM, Kopecky KJ, et al. Revised recommendations of the International Working Group for Diagnosis, Standardization of Response Criteria, Treatment Outcomes, and Reporting Standards for Therapeutic Trials in Acute Myeloid Leukemia. *J Clin Oncol*. 2003;21(24):4642-4649.
65. Cheson BD, Pfistner B, Juweid ME, et al. Revised response criteria for malignant lymphoma. *J Clin Oncol*. 2007;25(5):579-586.
66. Buchmann S, Schrappe M, Baruchel A, et al. Remission, treatment failure, and relapse in pediatric ALL: an international consensus of the Ponte-di-Legno Consortium. *Blood*. 2022;139(12):1785-1793.
67. Gökbuget N, Brüggemann M, Beck J, et al. Evaluation of minimal residual disease (MRD) and MRD-based treatment decisions in Ph/BCR-ABL negative adult acute lymphoblastic leukemia (ALL): experience from the German multicenter study group for adult ALL (GMALL). *Blood*. 2017;130:138.
68. Kotrova M, Koopmann J, Trautmann H, et al. Prognostic value of low-level MRD in adult acute lymphoblastic leukemia detected by low- and high-throughput methods. *Blood Adv*. 2022;6(10):3006-3010.
69. Cheson BD, Fisher RI, Barrington SF, et al. Recommendations for initial evaluation, staging, and response assessment of Hodgkin and non-Hodgkin lymphoma: the Lugano classification. *J Clin Oncol*. 2014;32(27):3059-3068.
70. Dreyling M, Andre M, Gokbuget N, et al. The EHA research roadmap: malignant lymphoid diseases. *Hemasphere*. 2022;6(6):e726.
71. Dohner H, Estey EH, Amadori S, et al. Diagnosis and management of acute myeloid leukemia in adults: recommendations from an international expert panel, on behalf of the European LeukemiaNet. *Blood*. 2010;115(3):453-474.

© 2024 American Society of Hematology. Published by Elsevier Inc. All rights are reserved, including those for text and data mining, AI training, and similar technologies.