



**1st
European
Conference on
Infection in
Leukemia**

EMPIRICAL ANTIBACTERIAL TREATMENT:
GLYCOPEPTIDES AND OTHER GRAM-
POSITIVE ANTIBACTERIALS

A.COMETTA, O.MARCHETTI, T.CALANDRA


Sept. 30th / Oct. 1st 2005 Juan-les-Pins - France



BACKGROUND

1. **Epidemiological data in the mid 80'**
2. **Development of resistance to glycopeptides in 90'**

IATG-EORTC TRIALS 1973-2000



1986-88

GLYCOPEPTIDES (GP) IN NEUTROPENIC PATIENTS: OBJECTIVES

- 1. Should GP be given as upfront empirical therapy ?**
- 2. Should GP be given in case of documented Gram positive MDI?**
- 3. Should GP be given in case of persistent fever after initial broad spectrum empirical antibiotic therapy?**



GLYCOPEPTIDES IN NEUTROPENIC PATIENTS: METHODS

- **Literature review**
 - **Search**
 - **Medline**
 - **Cochrane**
 - **Pubmed**
 - **Manual search bibliography of referenced publications**
 - **ICAAC, ECCMID, ASH, ASCO, and EBMT 2002-2005**
- **CDC grading**
- **Questionnaire on European practices.**

GLYCOPEPTIDES IN NEUTROPENIC PATIENTS: METHODS

- 1. Randomized controlled trials**
- 2. Meta-analysis**
 - 1. Paul et al JAC 2005; 55: 436-444**
 - 2. Vardakas Lancet Infect Dis 2005; 5: 431-439**
- 3. Published guidelines**

GLYCOPEPTIDES IN NEUTROPENIC PATIENTS

- 1. Upfront empirical therapy**
- 2. In case of persistent fever after initial broad spectrum empirical antibiotic therapy**
- 3. In case of documented Gram positive MDI**

RANDOMIZED CONTROLLED TRIALS WITH THE SAME ANTIBIOTIC(S) IN THE 2 GROUPS (1)

Trial/year	N=	Antibiotic	Glycopeptide
Karp 1986	60	Ticar-genta	Vancomycin
Del Favero 1987	47	Cefta-amika	Teicoplanin
Micozzi 1990	46	Pipera-amika	Teicoplanin
De Pauw 1990	103	Cefta	Teicoplanin
EORTC 1991	747	Cefta-amika	Vancomycin

RANDOMIZED CONTROLLED TRIALS WITH THE SAME ANTIBIOTIC(S) IN THE 2 GROUPS (2)

Trial/year	N=	Antibiotic	Glycopeptide
Novakova 1991	103	Cefta	Vancomycin
Ramphal 1992	127	Cefta	Vancomycin
Martino 1992	158	Pipera-amika	Teicoplanin
Pico 1993	102	Cefta	Vancomycin

RANDOMIZED CONTROLLED TRIALS WITH DIFFERENT ANTIBIOTICS IN THE 2 GROUPS (1)

Trial/year	N=	Antibiotic- no GP	Antibiotic + GP
Shenep 1988	101	Ticar-amika	Ticar/clav-amika
Meunier 1990	75	Cefta-amika	Cefta
Viscoli 1991	193	Cefta-amika	Cefta
Riikonen 1991	89	Imipenem	Cefta
Bosseray 1992	87	Imipenem	Cefta

RANDOMIZED CONTROLLED TRIALS WITH DIFFERENT ANTIBIOTICS IN THE 2 GROUPS (2)

Trial/year	N=	Antibiotic-no GP	Antibiotic + GP
Spencer 1990	59	Pip-genta	Aztreonam
Kelsey 1992	71	Pip-genta	Cefta
Micozzi 1993	104	Pip-amika	Pip/tazo-amika
Granowetter 1988	151	Carbeni-cephalo-genta	cefta

GLYCOPEPTIDES AS UPFRONT THERAPY

- 1. Mortality**
- 2. Success, duration of fever, shock**
- 3. Further infections, breakthrough bacteremia**
- 4. Toxicity**

Reference

Micozzi⁷

Pico⁸

Kelsey¹⁰

Martino¹¹

Ramphal¹²


Novakova¹⁴

Viscoli¹⁶

Meunier¹⁷

Shenep¹⁸

Combined



1. Odds ratios of mortality

Vardakas Lancet Infect Dis 2005; 5: 431-439



Infection
Leukemia

MORTALITY (1)

Trial/year	No Glycopeptide Death/total	Glycopeptide Death/total
Micozzi 1993	3/56	3/58
Kelsey 1992	2/29	1/29
Martino 1992	4/83	5/75
Ramphal 1992	6/63	7/64
Novakova 1991	9/60	7/60
Meunier 1990	9/50	8/50
Shenep 1988	1/48	0/53



MORTALITY (2)

Trial/year	No Glycopeptide Death/total	Glycopeptide Death/total
De Pauw 1990	6/51	4/52
EORTC 1991	19/370	24/377
Viscoli 1991	7/95	2/98
Pico 1993	10/69*	0/33


*** Ceftazidime 1g q 8h**

GLYCOPEPTIDES AS UPFRONT THERAPY

1. **Mortality**
2. **Shock, success, duration of fever**
3. **Further infections, breakthrough bacteremia**
4. **Toxicity**



2. Odds ratios of success without modification



Vardakas Lancet Infect Dis 2005; 5: 431-439

Initial addition of vancomycin for the empirical treatment of Gram-positive bacteremia in neutropenic patients

Modification of initial empirical treatment	Cefta-amika (n = 68)	Cefta-amika + vancomycin (n = 67)	
Vancomycin	22%	0%	<0.001
Other antibiotic	10%	12%	
Amphotericin B	10%	21%	<0.001
Acyclovir	8%	11%	

2. Time to defervescence

- **EORTC : no difference**
- **Karp: significant difference (median 14 days in placebo group vs 9 days in GP group)**
- **Meta-analysis: pooling data from 2 trials: no difference**

3.BREAKTHROUGH INFECTION (1)

Trial/year	No Glycopeptide /total	Glycopeptide /total
EORTC 1991	50/370 (13.5%)	42/377 (11%)
Novakova 1991	6/51	8/52
Viscoli 1991	9/63	11/75
Kelsey 1992	2/35	3/36
Ramphal 1992	8/63	5/64
Micozzi 1993	9/58	7/56
Bosseray 1992	1/43	1/44

3. BREAKTHROUGH INFECTION (2)

Trial/year	No Glycopeptide /total	Glycopeptide /total
Karp 1986	7 (32%)*	0
Marie/Pico 1993	35/146 (24%) G+ : 29/146	5/77 (6.5%) G+: 2/77

* Late onset G+ sepsis

3. G+ BREAKTHROUGH BACTEREMIA



Trial/year	No Glycopeptide n/total	Glycopeptide n/total
Shenep 1988	9/48*	1/53
Riikonen 1991	1/45	0/44
Granowetter 1988	1/55	1/46
Kelsey 1990	0/35	1/38

* CNS: 5. Viridans streptococci: 4 (1 death due to shock)

4. Odds ratio of adverse effects

A. All adverse effects

B. nephrotoxicity



4. ADVERSE EFFECTS (1)

Trial/year	No Glycopeptide n/total	Glycopeptide n/total
Bosseray 1992	0/43	2/44
Kelsey 1992	8/35	8/36
Martino 1992	0/83	2/75
Ramphal 1992	6/63	19/64
Viscoli 1991	4/95	34/98
Riikonen 1991	0/45	3/44
Del Favero 1987	4/33	6/33



4. ADVERSE EFFECTS (2) EORTC 1991

Adverse effect	No Glycopeptide n=370	Glycopeptide n= 383
Nephrotoxicity	9 (2%)	24 (6%)
Hepatotoxicity	50 (13.5%)	85 (22%)
Hypokaliemia	35 (9%)	55 (14%)
Rash	12 (3%)	26 (7%)

EORTC-IATCG, J Infect Dis, 1991; 163: 951-958

4. ADVERSE EFFECTS (3): nephrotoxicity

Trial/year	No Glycopeptide /total	Glycopeptide /total
Karp 1986	23/29	22/31
Kelsey 1992	1/35	0/36
Martino 1992	0/83	0/75
Riikonen 1991	0/45	0/44
Del Favero 1987	0/33	0/33
Novakova 1991	3/51	4/52
Meunier 1990	0/36	3/39



GLYCOPEPTIDES IN NEUTROPENIC PATIENTS

- 1. Upfront empirical therapy**
- 2. In case of documented Gram positive MDI**
- 3. In case of persistent fever after initial broad spectrum empirical antibiotic therapy**




Bacteremia due to viridans streptococci in granulocytopenic cancer patients

Trial/year	Pts number	Bacteremia due to S.viridans
Feld 2000	409	19 (4.6%)
Del Favero 2001	733	31 (4.3%)
Fleishback 2001	342	10 (2.9%)
Cordonnier 2003	513	24 (4.6%)
IATG-EORTC 2003	763	36 (4.7%)

EORTC-IATCG trial V: Gram-positive bacteremias

	Ceftazidime + Amikacin (n=68)	Ceftazidime + amikacin + vancomycin (n=67)
Streptococci	30	27
viridans	21	23
Coagulase-neg. staph.	28	21
<i>S. aureus</i>	4	16
Other	6	3

Initial addition of vancomycin for the empirical treatment of Gram-positive bacteremia in neutropenic patients



1st
European
Conference
Infection in
Leukemia

PATIENTS WITH SKIN AND SOFT TISSUE INFECTIONS

	Mono N=367	Comb N=355	Mono + V N=53	Comb + V N=43
Success (%)	35	33	42	42
Infectious mortality (%)	6	8	6	7
Days to defervescence	7.6	7.5	7.7	8.0
Superinfection (%)	10	10	15	8

GLYCOPEPTIDES IN NEUTROPENIC PATIENTS

1. **Upfront empirical therapy**
2. **In case of documented Gram positive MDI**
3. **In case of persistent fever after initial broad spectrum empirical antibiotic therapy:**
 - **Cometta et al CID 2003; 37: 382**
 - **Erjavec et al JAC 2000; 45: 843**

Addition of glycopeptides in neutropenic cancer patients

Trial/year	Pts number	pts with addition of glycopeptides
De Pauw 1994	722	26 %
IATG-GIMEMA 1996	987	36%
Winston 2001	541	31%
Sanz 2002	867	45%
Peacock 2002	471	62%

Day 0

859 febrile neutropenic Pts

**763 eligible pts:
piperacillin/tazobactam**

96 Pts not eligible

48-60 hours

**165 Pts with persistent
fever and FUO, CDI or
Bacteremia due to G+
susceptible to P/T**

**598 Pts afebrile, or with
exclusion criteria for
randomization**

RANDOMIZATION


STUDY OF P/T EFFICACY



Randomized patients: defervescence

	Placebo N = 79	Vancomycin N = 86
Pts with defervescence	73 (92%)	82 (96%)
Pts with defervescence under protocol therapy	36 (45%)	42 (49%)
Pts with defervescence after change of protocol therapy	37 (47%)	40 (47%)
Median time to defervescence (Days; 95% C.I.)	4.3 (3.3-4.7)	3.5 (2.7-4.4)

Overall time to defervescence



Infection in
Leukemia

Outcome of the patients

	Placebo N = 79	Vancomycin N = 86
Further G+ bacteremia	4	3
Pts given ampho B	30 (37%)	31 (36%)
Pts with AE definitely or probably related to AB	3	9
Death related to infection (Day of death)	2 (15, 35)	1 (14)

Day 0

X febrile neutropenic Pts

72-96 hours

**124 pts:
imipenem/cilastatin**

**11 Pts
not eligible**

**115 Pts with persistent
fever and FUO, CDI or
Bacteremia due to G+
susceptible to I/C**

RANDOMIZATION

Erjavec et al outcome of the patients

	Placebo N = 58	Teicoplanin N = 56
Pts with defervescence	27 (46.6%)	25 (44.6%)
Death	4 (6.9%)	6 (10.7%)

1. Initial empirical glycopeptide in neutropenic patients (IDSA 2002)

- **Development of hypotension or shock**
- **Known colonisation with MRSA or Peni-R Pneumococcus**
- **Positive results for G+ before identification**
- **Clinically suspected serious cath-related infection (cellulitis)**
- **(Institutions with high rate of infections due to MRSA or Peni-R viridans streptococci)**

RANDOMIZED CLINICAL TRIALS: PROBLEMS

- **No double-blind trial except Karp's and Shenep's trials: addition of GP more frequent in the group initially treated without GP**
- **More trials with different antibiotics in the 2 groups: role in the occurrence of adverse effects and further infections?**
- **Various doses of vancomycin and teicoplanin**
- **No randomized controlled trial assessing the use of streptogramin or linezolid**

CONCLUSION 1

	Glycopeptide	CDC grading system
At onset of fever	Not recommended	I D
Persistent fever	Not recommended	I D

CONCLUSION 2

	Glycopeptide	CDC grading system
Known colonisation with MRSA	recommended	III C
Hypotension or shock	recommended	III C
Skin and soft tissue infections including cath-related infections	recommended	III C