

Öffentlicher Titel	Phase II/III Studie zu Imetelstat bei myelodysplastischen Syndromen mit niedrigem oder mittlerem IPSS-Risiko
Wissenschaftl. Titel	A Study to Evaluate Imetelstat (GRN163L) in Transfusion-Dependent Subjects with IPSS Low or Intermediate-1 Risk Myelodysplastic Syndrome (MDS) that is Relapsed/Refractory to Erythropoiesis-Stimulating Agent (ESA) Treatment
Kurztitel	63935937MDS3001
Studiennummer KN/ELN	LN_NN_2020_697
Studiengruppe	NN
Studienart	randomisiert, prospektiv, doppelblind
Studienphase	Phase II/III
Erkrankung	Myelodysplastisches Syndrom (MDS) - Niedrigrisiko und Intermediär I
Einschlusskriterien	<ul style="list-style-type: none">- Man or woman greater than or equal to (\geq) 18 years of age- In Part 1, diagnosis of myelodysplastic syndrome (MDS) according to World Health Organization (WHO) criteria- International Prognostic Scoring System (IPSS) low Risk or intermediate-1 risk MDS- Red blood cell (RBC) transfusion dependent, defined as requiring at least 4 RBC units transfused over an 8-week period during the 16 weeks prior to Study Entry; pre-transfusion hemoglobin (Hb) should be less than or equal to 9.0 gram per deciliter (g/dL) to count towards the 4 units total- Eastern Cooperative Oncology Group (ECOG) performance status 0, 1 or 2
Ausschlusskriterien	<ul style="list-style-type: none">- Participant has known allergies, hypersensitivity, or intolerance to imetelstat or its excipients- Participant has received an investigational drug or used an invasive investigational medical device within 30 days prior to Study Entry or is currently enrolled in an investigational study- Prior treatment with imetelstat- Have received corticosteroids greater than ($>$) 30 milligram per day (mg/day) prednisone or equivalent, or growth factor treatment within 4 weeks prior to study entry- a) Prior treatment with a hypomethylating agent (example [eg], azacitidine, decitabine); b) Prior treatment with lenalidomide; c) Has received an erythropoiesis-stimulating agent (ESA) or any chemotherapy, immunomodulatory, or immunosuppressive therapy within 4 weeks prior to study entry (8 weeks for long-acting ESAs)
Alter	\geq 18 Jahre
Status	Aktiv
Beginn der Rekrutierung	10.03.2020
Sponsoren	Geron Corporation
Registrierung in anderen Studienregistern	ClinicalTrials.govNCT02598661 European Clinical Trials Database - EUDRACT201500287419